

Monitoring Cell Migration

Jeff W.M. Bulte¹ and Michel M. Modo²

¹*Russell H. Morgan Dept. of Radiology and Radiological Science, Div. of MR Research, and Institute for Cell Engineering, Johns Hopkins Univ. School of Medicine, Baltimore, MD;*

²*Neuroimaging Research Group and Center for the Cellular Basis of Behaviour, Institute of Psychiatry, King's College London, London, UK .*

MR cellular imaging is a relatively new and emerging technique aimed at guiding development of effective cell therapies, including progenitor cells, stem cells, and immunotherapeutic cells. It provides the opportunity to non-invasively map grafted cells in living animals and humans, to observe their migration and, more importantly, to link functional measures with the presence of the cells in target organ. This technique also addresses the safety issue by monitoring potential uncontrolled proliferation or dispersion to unwanted areas. At the present time, MR cellular imaging now allows whole-body scanning capability at a resolution approaching the size of individual transplanted cells (1, 2), while allowing repeated analysis at multiple time points.

Numerous techniques have been developed to magnetically label cells prior to grafting, most of them utilizing dextran-coated superparamagnetic iron oxide nanoparticles (SPIOs). SPIOs can be used to label cells from multiple species, simply by 24-48 hours of incubation in conjunction with transfection agents (3) or, as has been recently shown, by instant labeling using magnetoelectroporation (4), allowing for non-invasive detection of transplants using MRI. Most recent studies have employed Feridex as SPIO, as it is a commercially-available and FDA-approved source of dextran-coated SPIO. Gd-chelates have also been used (5-7), albeit in much less extent, owing to the reduced sensitivity of the paramagnetic label once internalized in cells, forcing the use of T2(*)-weighted hypointensity imaging at higher field strengths. A large number of gadolinium molecules are needed in each cell to allow a robust *in vivo* visualization, although some strategies, such as metallofullerenes (8), have been described to increase relaxivity and therefore could reduce the number of paramagnetic molecules needed for cellular imaging. Daldrup-Link (9) demonstrated that twice as many cells labeled with gadopentate dimeglumine liposomes were needed for detection by MRI compared to ferumoxides, P7728 liposomes or polysaccharide nanoparticles.

Most interest to date in the use of pre-labeled cells for cellular MR imaging has been in the use of stem cells. The *ex vivo* MR analysis of the global or local distribution of transplanted stem cells is now a routinely used method that complements conventional histological techniques. This method allows for 3D whole body or organ imaging and presentation of data in any desired anatomical plane—features that are impossible or extremely difficult to achieve using conventional microscopy. Post-mortem (*ex vivo*) imaging is not subject to the MR technique limitations associated with living organism imaging (limited acquisition time or motion artifacts); thus, the spatial resolution, signal-to-noise ratio, and sensitivity are much better. The MRI cell tracking technique, however, offers far more valuable data when used in a living organism, allowing for dynamic evaluation of cell migration. From a clinical perspective, it also offers a non-invasive means of monitoring cell-based therapy that enables determining the cell

fate without the need for tissue biopsy. Brain and cardiac disorders have received so far the most attention for the potential application of MRI stem cell tracking techniques (10).

Monitoring immune cell trafficking is the second major application of MR imaging of cell migration. It is likely to have a significant future impact on our understanding of the *in vivo* immune response. Anderson et al. have demonstrated targeted trafficking of myelin—sensitized T cells into the spinal cord of animals with experimental allergic encephalomyelitis (11). Antigen-specific cytotoxic T cells have been found to specifically home to OVA⁺ (12) and HER2/neu tumors (13). T cells from animals that rejected OVA⁺ tumors have been purified, SPIO-labeled, and re-injected in OVA⁺ tumor-bearing animals (14). In this case, specific tumor accumulation could also be observed. MR monitoring of cytotoxic T cell migration into the diabetic pancreas has been pursued in order to gain insight into the mechanisms underlying autoimmune diabetes (15, 16). SPIO-labeled cells behave clearly different from naked SPIO particles when injected intravenously; Endorem-labeled hematopoietic bone marrow cells were found to rapidly home to bone marrow following intravenous injection whereas just the Endorem particles by themselves do not (17).

While MR imaging is rapidly gaining ground for cellular imaging, other techniques also possess the potential for cellular imaging in the living organism. Especially studies on hematopoietic stem cell transplantation have used fluorescent dyes, such as PKH26, to pre-label cells *in vitro* to allow *in vivo* detection of transplanted/infused cells by means of optical imaging (18-20). However, this approach does not allow deep tissue penetration and therefore is limited to easily accessible targets. Alternatively, invasive procedures, such as a craniectomy, are required to allow access of the imaging probe to the anatomical structure of interest (18). Exposure to light to visualize the fluorescent dyes can also cause phototoxicity (21) killing the labeled cells complicating the serial assessment of these cells by optical imaging methods. The attractiveness of this approach lies in the commercial availability of fluorescent dyes for cellular tracking that are widely used on a variety of cell types and provides a powerful approach to study, for instance, cellular differentiation by combining the fluorescent marker for the identification of transplanted cells with immunohistochemical methods.

A promising new development in cellular imaging consists of the possibility to detect bioluminescence from deep tissue structures (22). Typically, a gene encoding for luciferase, is engineered into cells (23). The substrate luciferin (supplied through i.v. injection) is oxidized by luciferase in the presence of ATP and oxygen, producing a photon in the course of this bioluminescent reaction that can be detected by very sensitive cameras, thus allowing repeated *in vivo* cellular imaging (24). A drawback of bioluminescent imaging is that the scattering of light by tissue limits the spatial resolution of this technique considerably compared to MRI. In addition, clinical or whole body large animal imaging is, at present, not feasible, and pigmented strains are less suitable for bioluminescent imaging, raising ethical issues would the technology allow imaging in humans. Still, bioluminescent imaging has allowed Kim and colleagues (25) to track the *in vivo* migration of neural stem cells from the contralateral hemisphere to the site of lesion in mice with middle cerebral artery occlusion whilst allowing a corroboration of these results by histology. Migration of neural stem cells into gliomas across hemispheres has also been visualized (26), and the trafficking of immune cells has been monitored as well (27, 28). The potential to track a particular type of cell, such as a neural stem cell or immune cell, by bioluminescence, whilst using the same techniques to detect light emitted at a different wavelength from distinct reporter genes to study gene expression in these cells clearly will provide interesting novel avenues to study the basic biology of transplanted cells by combining

molecular (gene expression) and cellular imaging (location of cells).

Various other techniques, such as positron emission tomography (PET) and single photon emission tomography (SPECT), also have the potential to visualize cells and cell migration *in vivo*. The current spatial resolution of both PET and SPECT is at least 10 times lower than that observed with MRI, but it is the higher specificity of radioligands that conveys the attractiveness of these techniques to study transplanted cells *in vivo* (29). For instance, Jacobs et al (30) have shown that engineering of reporter genes into cells *in vitro* allows the identification of these cells by means of detection of the transgene expression *in vivo*. Hence, the reporter molecules could serve as a surrogate marker of the cells to be imaged and no prior loading of the cells with a contrast agent would be required as radioligands are small enough to cross the intact blood-brain barrier. Differential expression of proteins/receptors, such as the transferrin receptor, on particular cells can also serve as a marker to identify these transplanted cells *in vivo* (31, 32). Although both PET and SPECT provide excellent approaches to study specific molecular targets and determine cellular differentiation or metabolic consumption, the repeated assessment of animals with radioligands raises issues regarding the safety of this approach for longitudinal studies and its translation into a clinical setting. The pre-labeling of cells *in vitro* with radioligands prior to *in vivo* imaging (as exemplified by MR cellular imaging) is also less likely to be apposite for the longitudinal tracking of cells due to the short half-life of radioligands. Nonetheless, Chin et al. (33) have demonstrated that it is possible to visualize mesenchymal stem cells by SPECT over a period of 14 days after pre-incubation of cells with ^{111}In oxine. Kraitchman et al. (34) clearly demonstrated the much higher sensitivity of detecting mesenchymal stem cells with ^{111}In oxine as compared to labeling with Feridex; estimated numbers of between 50,000-100,000 cells could be detected as hot spots on SPECT/CT images but not on the MR images (using clinically applicable imaging parameters). Another cell tracking approach is the use of specific reporter markers in transplanted cells that are detectable by repeated administration of a radioligand thus providing an attractive approach to monitor immune (35, 36) or stem cell (37) graft survival and function over extended time periods.

However, the need for on-site synthesis of radioligands limits the wider applicability and practicality of PET and SPECT as cellular imaging techniques, whereas limits on tissue penetration and need for genetic engineering will complicate translation of optical imaging and bioluminescence into clinical applications. With MR imaging, it is impossible to differentiate live from dead cells, and label dilution by cell division is a significant limitation. But the wider availability of MRI scanners in both clinical and pre-clinical settings strengthens the applicability of cellular MR imaging as a translational technique that can be easily implemented in many centers. Given the high-resolution of MR imaging at near cellular levels, however, it is the unique complementary information that can be derived from the different imaging approaches that highlight their importance in cellular imaging. This is well illustrated by the first clinical trial that has used magnetically labeled cells (38): MR imaging was superior to ^{111}In -oxine scintigraphy in detecting the number of lymph nodes that Endorem-labeled dendritic cells migrated to following intranodal injection in a single regional node, and was able to accurately assess whether the nodes were hit or missed following US-guided injections. At the other hand, only scintigraphy allowed accurate quantification of migrated cell numbers, which is important for predicting sufficient T cell activation.

For cellular MR imaging, a few new approaches were introduced last year that may offer opportunities for improved monitoring of cell migration. Gilad et al. in our group have developed an MR reporter gene, that is unique in the sense that it is an artificial reporter gene providing

endogenous contrast without the need for administration of contrast agents (39). Using chemical exchange saturation transfer (CEST) imaging, images can be created using specific off-radiation pulses that null the signal of amide protons in the reporter gene. By subtracting and correcting the anatomical images without the specific pulse, “hot spot” images can be created that have now allowed accurate detection of cell distribution in vivo. Successful development of MR reporter genes would have the same advantages that bioluminescent and PET imaging have when using reporter genes: 1) no decrease of the sensitivity of detection following cell division (caused by dilution of label); 2) the ability to discriminate between live and dead cells (as only live cells continue to produce the reporter); and 3) the potential to place reporters under specific promoters thus allowing the detection of cells only under certain activation or differentiation conditions (for instance, the visualization of mesenchymal stem cells only when differentiated into cardiomyocytes). Cohen et al. (40) Genove et al. (41) have used a different approach that still relies on metal-based relaxation changes, namely the use of a ferritin reporter gene, synthesizing its own iron oxyhydroxide core, that could possibly also be used for cell tracking.

Finally, ^{19}F -based “hot spot” MR imaging has emerged from the background. Perhaps largely unrecognized, fluorine-19 was one of the earliest applied MR contrast agents (42). Ahrens et al. (43) have used large particles containing many ^{19}F atoms to label dendritic cells, and demonstrated that organ trapping in the liver, spleen and lungs could be readily visualized. As compared to ^1H imaging and using metal-based contrast agents, ^{19}F -based imaging offers several advantages but may also have some pitfalls, which have recently been discussed in further detail (44). Thus, there are many ways to monitor cell migration. It is too early to tell which of all the described techniques will become the future technique of choice, but it is clear that there is an array of possibilities with the optimal choice depending on the specific cell tracking application.

References

1. Shapiro EM, Sharer K, Skrtic S, Koretsky AP. In vivo detection of single cells by MRI. *Proc. Int. Soc. Magn. Reson. Med.* 2005; 13:355.
2. Heyn C, Ronald JA, Mackenzie LT, et al. In vivo magnetic resonance imaging of single cells in mouse brain with optical validation. *Magn. Reson. Med.* 2006; online in advance of print.
3. Frank JA, Miller BR, Arbab AS, et al. Clinically Applicable Labeling of Mammalian and Stem Cells by Combining Superparamagnetic Iron Oxides and Transfection Agents. *Radiology* 2003; 228:480-487.
4. Walczak P, Kedziorek DA, Gilad AA, Lin S, Bulte JW. Instant MR labeling of stem cells using magnetoelectroporation. *Magn Reson Med* 2005; 54:769-774.
5. Modo M, Cash D, Mellodew K, et al. Tracking transplanted stem cell migration using bifunctional, contrast agent-enhanced, magnetic resonance imaging. *Neuroimage* 2002; 17:803-811.
6. Crich SG, Biancone L, Cantaluppi V, et al. Improved route for the visualization of stem cells labeled with a Gd-/Eu-chelate as dual (MRI and fluorescence) agent. *Magnetic Resonance in Medicine* 2004; 51:938-944.
7. Zheng Q, Dai H, Merritt ME, Malloy C, Pan CY, Li WH. A new class of macrocyclic lanthanide complexes for cell labeling and magnetic resonance imaging applications. *J Am Chem Soc* 2005; 127:16178-16188.
8. Mikawa M, Kato H, Okumura M, et al. Paramagnetic water-soluble metallofullerenes having the highest relaxivity for MRI contrast agents. *Bioconjugate Chemistry* 2001; 12:510-514.
9. Daldrop-Link HE, Rudelius M, Oostendorp RA, et al. Targeting of hematopoietic progenitor cells with MR contrast agents. *Radiology* 2003; 228:760-767.
10. Bulte JW, Kraitchman DL. Monitoring cell therapy using iron oxide MR contrast agents. *Curr Pharm Biotechnol* 2004; 5:567-584.
11. Anderson SA, Shukaliak-Quandt J, Jordan EK, et al. Magnetic resonance imaging of labeled T-cells in a mouse model of multiple sclerosis. *Ann Neurol* 2004; 55:654-659.

12. Kircher MF, Allport JR, Graves EE, et al. In vivo high resolution three-dimensional imaging of antigen-specific cytotoxic T-lymphocyte trafficking to tumors. *Cancer Res* 2003; 63:6838-6846.
13. Daldrup-Link HE, Meier R, Rudelius M, et al. In vivo tracking of genetically engineered, anti-HER2/neu directed natural killer cells to HER2/neu positive mammary tumors with magnetic resonance imaging. *Eur Radiol* 2005; 15:4-13.
14. Hu DE, Kettunen MI, Brindle KM. Monitoring T-lymphocyte trafficking in tumors undergoing immune rejection. *Magn Reson Med* 2005; 54:1473-1479.
15. Moore A, Sun PZ, Cory D, Hogemann D, Weissleder R, Lipes MA. MRI of insulinitis in autoimmune diabetes. *Magn Reson Med* 2002; 47:751-758.
16. Billotey C, Aspod C, Beuf O, et al. T-cell homing to the pancreas in autoimmune mouse models of diabetes: in vivo MR imaging. *Radiology* 2005; 236:579-587.
17. Daldrup-Link HE, Rudelius M, Piontek G, et al. Migration of iron oxide-labeled human hematopoietic progenitor cells in a mouse model: in vivo monitoring with 1.5-T MR imaging equipment. *Radiology* 2005; 234:197-205.
18. Askensay N, Farkas DL. Optical imaging of PKH-labeled hematopoietic cells in recipient bone marrow in vivo. *Stem Cells* 2002; 20:501-513.
19. Fox D, Kouris GJ, Blumofe KA, Heilizer TJ, Husak V, Greisler HP. Optimizing fluorescent labeling of endothelial cells for tracking during long-term studies of autologous transplantation. *Journal of Surgical Research* 1999; 86:9-16.
20. Ferrari A, Hannouche D, Oudina K, et al. In vivo tracking of bone marrow fibroblasts with fluorescent carbocyanine dye. *Journal of Biomedicine and Material research* 2001; 56:361-367.
21. Oh DJ, Lee GM, Francis K, Palsson BO. Phototoxicity of the fluorescent membrane dyes PKH2 and PKH26 on the human hematopoietic KG1a progenitor cell line. *Cytometry* 1999; 36:312-318.
22. Contag CH, Bachmann MH. Advances in in vivo bioluminescence imaging of gene expression. *Annual Reviews in Biomedical Engineering* 2002; 4:235-260.
23. Rice BW, Cable MD, Nelson MB. In vivo imaging of light-emitting probes. *J Biomed Opt* 2001; 6:432-440.
24. Koransky ML, Ip TK, Wu S, et al. In vivo monitoring of myoblast transplantation into rat myocardium. *J Heart Lung Transplant* 2001; 20:188-189.
25. Kim D-E, Schellingerhout D, Ishii K, Shah K, Weissleder R. Imaging of stem cell recruitment to ischemic infarcts in a murine model. *Stroke* 2004; 35:952-957.
26. Tang Y, Shah K, Messerli SM, Snyder E, Breakefield X, Weissleder R. In vivo tracking of neural progenitor cell migration to glioblastomas. *Hum Gene Ther* 2003; 14:1247-1254.
27. Costa GL, Sandora MR, Nakajima A, et al. Adoptive immunotherapy of experimental autoimmune encephalomyelitis via T cell delivery of the IL-12 p40 subunit. *J Immunol* 2001; 167:2379-2387.
28. Hardy J, Edinger M, Bachmann MH, Negrin RS, Fathman CG, Contag CH. Bioluminescence imaging of lymphocyte trafficking in vivo. *Exp Hematol* 2001; 29:1353-1360.
29. Marzullo P. Nuclear imaging after cell implantation. *Int J Cardiol* 2004; 95 Suppl 1:S53-54.
30. Jacobs A, Tjuvajev JG, Dubrovin M, et al. Positron emission tomography-based imaging of transgene expression mediated by replication-conditional, oncolytic herpes simplex virus type 1 mutant vectors in vivo. *Cancer Res* 2001; 61:2983-2995.
31. Ding W, Bai J, Zhang J, et al. In vivo tracking of implanted stem cells using radio-labeled transferrin scintigraphy. *Nucl Med Biol* 2004; 31:719-725.
32. Bai J, Ding W, Yu M, et al. Radionuclide imaging of mesenchymal stem cells transplanted into spinal cord. *Neuroreport* 2004; 15:1117-1120.
33. Chin BB, Nakamoto Y, Bulte JWM, Pittenger MF, Wahl R, Kraitchman DL. ¹¹¹In oxine labelled mesenchymal stem cell SPECT after intravenous administration in myocardial infarction. *Nuclear Medicine Communications* 2003; 24:1149-1154.
34. Kraitchman DL, Tatsumi M, Gilson WD, et al. Dynamic imaging of allogeneic mesenchymal stem cells trafficking to myocardial infarction. *Circulation* 2005; 112:1451-1461.
35. Koehne G, Dubrovin M, Dubrovina E, et al. Serial in vivo imaging of the targeted migration of human HSV-TK-transduced antigen-specific lymphocytes. *Nature Biotechnology* 2003; 21:405-413.
36. Dubey P, Su H, Adonai N, et al. Quantitative imaging of the T cell antitumor response by positron-emission tomography. *Proc Natl Acad Sci U S A* 2003; 100:1232-1237.
37. Wu JC, Chen IY, Sundaresan G, et al. Molecular imaging of cardiac cell transplantation in living animals using optical bioluminescence and positron emission tomography. *Circulation* 2003; 108:1302-1305.

38. de Vries IJ, Lesterhuis WJ, Barentsz JO, et al. Magnetic resonance tracking of dendritic cells in melanoma patients for monitoring of cellular therapy. *Nat Biotechnol* 2005; 23:1407-1413.
39. Gilad AA, McMahon MT, Winnard Jr. PT, Raman V, Bulte JWM, van Zijl PCM. MRI reporter gene providing contrast based on chemical exchange saturation transfer (CEST). *Proc. Int. Soc. Magn. Reson. Med.* 2005; 13:363.
40. Cohen B, Dafni H, Meir G, Harmelin A, Neeman M. Ferritin as an endogenous MRI reporter for noninvasive imaging of gene expression in C6 glioma tumors. *Neoplasia* 2005; 7:109-117.
41. Genove G, DeMarco U, Xu H, Goins WF, Ahrens ET. A new transgene reporter for in vivo magnetic resonance imaging. *Nat Med* 2005; 11:450-454.
42. Holland GN, Bottomley PA, Hinshaw WS. ¹⁹F magnetic resonance imaging. *J. Magn. Reson.* 1977; 28:133-136.
43. Ahrens ET, Flores R, Xu H, Morel PA. In vivo imaging platform for tracking immunotherapeutic cells. *Nat Biotechnol* 2005; 23:983-987.
44. Bulte JW. Hot spot MRI emerges from the background. *Nat Biotechnol* 2005; 23:945-946.